Edited by: Edward T. Olsen

Contributions by:

J. Susskind, J. Blaisdell Goddard Space Flight Center, NASA



June, 2007 Version 1.0



Jet Propulsion Laboratory California Institute of Technology Pasadena, CA

Submit Questions to:

http://airs.jpl.nasa.gov/AskAirs

Table of Contents

TABLE OF CONTENTS	
INTRODUCTION TO V5 RETRIEVAL FLOW	
COMPARISON OF V4 AND V5 RETRIEVAL FLOWS	
NOTATION	6
Atmospheric States	6
Operations	6
Physical Parameters	7

INTRODUCTION TO V5 RETRIEVAL FLOW

All quantities marked **best** (associated quality factor = 0) and almost all quantities marked **good** (associated quality factor = 1) have passed through the entire retrieval system. It may be helpful to some users to understand the processing paths by which termination of the V5 retrieval process can occur. The steps are single-path sequential for V5, but early termination occurs when something fails along the way. There are seven possibilities, listed below.

- Retrievals falling into case 7 are considered to be successful MW/IR retrievals. Products should be used or avoided as indicated by their individual quality flags. The user is encouraged to read
 V5_L2_Quality_Control_and_Error_Estimation.pdf, which describes the individual quality flags.
- Retrievals falling into case 1 should not be used for any purpose.
- Retrievals falling into case 3 contain only microwave products.
- We recommend using only cloud products and OLR products from retrievals falling into cases 2, 4, 5 and 6, along with the cloud products falling into case 7.
 - MW retrieval fails (i.e., temperature profile is rejected), cloudy regression succeeds (or fails), and the first cloud parameter retrieval fails. All output is considered **bad** (associated quality factors set to 2) and most products contain only flag values. We estimate 1% of retrievals are in this subset.
 - 2. MW retrieval fails, but cloudy regression succeeds and first cloud parameter retrieval succeeds. In this case we attempt to continue only through this first cloud parameter retrieval and then terminate (there is no trusted MW emissivity.) Clouds and OLR are output with associated quality flags set to 1; all other quantities are output with associated quality flags set to 2. We estimate 1% of retrievals are in this subset.
 - 3. MW retrieval succeeds, but failure occurs before the end of the first cloud parameter retrieval. MW products are marked as described in V5_L2_Standard_Product_QuickStart.pdf. All other products are unusable and marked bad (associated quality flags set to 2). We estimate 1% of retrievals are in this subset.
 - 4. MW retrieval succeeds, first cloud parameter retrieval succeeds, but failure occurs before final temperature retrieval. The state at the end of the first cloud parameter retrieval (the "fallback state") is output, with clouds and OLR associated quality flags set to 1 and all other parameters associated quality flags set to 2. We estimate 2% of retrievals are in this subset.
 - 5. All steps succeed through final temperature retrieval, but failure occurs in CO or CH4 retrieval. CO and CH4 are marked bad (associated quality flags set to 2), but other states are marked as in (6) or (7) below. This situation is not known to occur.

- 6. All steps execute, but the case is determined to be at least 90% cloudy. The final retrieval is judged to be inferior to the state at the end of the first cloud parameter retrieval, and the fallback state is output. The quality factors for cloud products and OLR are set to 1 and all other quality factors set to 2. We estimate 5% of retrievals are in this subset.
- 7. All steps execute and the case is determined to be less than 90% cloudy. The quality factors for cloud products and OLR are set to 0. Quality control is applied to other parameters as described in the document, V5_L2_Quality_Control_and_Error_Estimation.pdf. We estimate 90% of retrievals are in this subset.

A step-by-step comparison of V4 and V5 retrieval flow follows.

COMPARISON OF V4 AND V5 RETRIEVAL FLOWS

Version 4	Version 5	Ver 5 AIRS-Only
$\mathbf{X}^{\mathbf{CLIM}}$, $\mathbf{R}_{\mathbf{MW}} \Rightarrow \mathbf{X}^{\mathbf{MW}}$,	\mathbf{X}^{CLIM} , $\mathbf{R}_{MW} \Rightarrow \mathbf{X}^{MW}$,	
$(\epsilon_{MW}, \mathbf{q}_{liq}, \mathbf{C}_{s})$	$(\epsilon_{MW}$, \mathbf{q}_{liq} , $\mathbf{C}_{s})$	
N/A	Cloudy Regression R_{MW} , $R_{IR} \Rightarrow X^{CR}$	Cloudy Regression $R_{IR} \Rightarrow X^{CR}$
AMSU Retrieval R _{MW} , X ^{MW} ⇒X ⁰	AMSU Retrieval R _{MW} , X ^{CR} ⇒X ⁰	set X ⁰ = X ^{CR}
(update T, T _s , q)	(update T , T _s , q)	
Cloud Parm Ret 1 α^0 , P_c^0	Cloud Parm Ret 1 α^0 , P_c^0	Cloud Parm Ret 1 α^0 , P_c^0
1 st Cloud Clearing R_{IR} , $X^0 \rightarrow \hat{R}_{IR}^0$	1 st Cloud Clearing R _{IR} , X ⁰ → R̂ ⁰ _{IR}	1 st Cloud Clearing R _{IR} , X ⁰ → R̂ _{IR} 0
Regression R̂ IR → XREG	Regression R̂ _{IR} →X ^{REG}	$ \begin{array}{c} \textbf{Regression} \\ \hat{\textbf{R}}_{\text{IR}}^{\text{0}} {\rightarrow} \textbf{X}^{\text{REG}} \end{array} $
$(T,T_{s},q,O_{3},\varepsilon_{IR})$	(T,T_s,q,ϵ_IR)	(T,T_s,q,ϵ_IR)
AMSU Retrieval R _{MW} , X ^{REG} ⇒X ¹ X ¹	AMSU Retrieval R _{MW} , X ^{REG} ⇒X ¹	
(update T , ε _{MW})	(update T , T_s , ϵ_{MW})	
Cloud Parm Ret 2 α^1 , P_c^1	Cloud Parm Ret 2 α^1 , P_c^1	Cloud Parm Ret 2 α^1 , P_c^1
(continued next page)	(continued next page)	(continued next page)

Version 4	Version 5	Ver 5 AIRS-Only
2 nd Cloud Clearing	2 nd Cloud Clearing	2 nd Cloud Clearing
R_{IR} , $X^1 \rightarrow \hat{R}_{IR}^1$	R_{IR} , $X^1 \rightarrow \hat{R}_{IR}^1$	R_{IR} , $X^{REG} \rightarrow \hat{R}_{IR}^1$
Physical Retrieval	NI/A	NI/A
R_{MW} , R_{IR} , $X^1 \Rightarrow T_s$	N/A	N/A
AMSU Retrieval		
R_{MW} , X^1 , $T_s \Rightarrow X^2$	N/A	N/A
(update T , ε _{MW})		
Cloud Parm Ret 3	N/A	N/A
α^2 , P_c^2	IN/A	11/7
3 rd Cloud Clearing	N/A	N/A
R_{IR} , $X^2 \rightarrow \hat{R}_{IR}^2$	IN/A	IN/A
Physical Retrieval	Physical Retrieval	Physical Retrieval
$X^2, \hat{R}_{IR}^2 \Rightarrow X^{PHYS}$	X^1 , $\hat{R}_{IR}^1 \Rightarrow X^{PHYS}$	$X^1, \hat{R}_{IR}^1 \Rightarrow X^{PHYS}$
$(T_s, \varepsilon_{IR}, T, q, O_3)$	$(T_s, \varepsilon_{IR}, T, q, O_3)$	$(T_{s}, \epsilon_{IR}, T, q, O_3)$
AMSU Retrieval	AMSU Retrieval	
R_{MW} , $X^{PHYS} \Rightarrow X^{test}$	R_{MW} , $X^{PHYS} \Rightarrow X^{test}$	N/A
(update T , ε _{MW})	(for quality control)	
Cloud Parm Ret 4	Cloud Parm Ret 3	Cloud Parm Ret 3
α^3 , P_c^3	α^2 , P_c^2	α^2 , P_c^2
4 th Cloud Clearing	3 rd Cloud Clearing	3 rd Cloud Clearing
R_{IR} , $X^{test} \rightarrow \hat{R}_{IR}^3$	R_{IR} , $X^{PHYS} \rightarrow \hat{R}_{IR}^2$	R_{IR} , $X^{PHYS} \rightarrow \hat{R}_{IR}^2$
Physical Retrieval	Physical Retrieval	Physical Retrieval
X^{test} , $\hat{R}_{IR}^3 \Rightarrow X^{\text{final}}$	$X^1, X^{PHYS}, \hat{R}_{IR}^2 \Rightarrow X^{final}$	$X^1, X^{PHYS}, \hat{R}_{IR}^2 \Rightarrow X^{final}$
$(T_s, \epsilon_{IR}, T, CO)$	$(T_s, \varepsilon_{IR}, T, CO, CH_4)$	$(T_s, \varepsilon_{IR}, T, CO, CH_4)$
Choose Final State	Choose Final State	Choose Final State
$X^{OUT} = X^0 \text{ or } X^{final}$	$X^{OUT} = X^0 \text{ or } X^{final}$	$X^{OUT} = X^0 \text{ or } X^{final}$
$X^{OUT} \rightarrow OLR$	X ^{OUT} → OLR	$X^{OUT} \rightarrow OLR$
	Error Estimates δX	Error Estimates δX
Quality Control	Quality Control	Quality Control

NOTATION

Atmospheric States

X^{CLIM} climatology atmospheric state, with the addition of the AVN surface pressure derived from the 3, 6 and 9 hour forecasts.

X^{MW} atmospheric state derived by MW-Only Retrieval (product)

X^{CR} atmospheric state derived by Cloudy Regression

X^{REG} atmospheric state derived by Regression

XPHY atmospheric state derived by Physical Retrieval

Xⁿ nth atmospheric state

X^{test} derived atmospheric state used only for quality control

X^{final} final atmospheric state derived by physical retrieval algorithm

X^{OUT} atmospheric state reported as product

δX error estimate for atmospheric state product

Note: The climatology atmospheric state is the NCEP 20-year temperature (T(p)) and moisture (q(p)) daily mean reanalysis on a 2-degree grid, extended above 100-mb by a UARS climatology and interpolated to the AIRS pressure levels.

Operations

X^a⇒X^b retrieval of atmospheric state "b" starting from state "a"

 $X^c \rightarrow \hat{R}_i^c$ derivation of parameter(s) from atmospheric state "c"

Physical Parameters

R_{MW} observed MW radiances

R_{IR} observed (cloudy) IR radiances

 $\hat{\mathbf{R}}_{IR}^{c}$ cloud cleared IR radiances derived from atmospheric state "c"

 α^n cloud fraction, iteration "n"

Pⁿ cloud top pressure, iteration "n"

q_{liq}, liquid water content from MW-Only Retrieval

C_s surface classification from MW-Only Retrieval

 ϵ_{MW} surface emissivity from MW-Only Retrieval

T_s surface skin temperature

OLR outgoing longwave radiation product

T Tair profile product

q precipitable water vapor profile/burden product

O₃ ozone profile/burden product

CO carbon monoxide profile/burden product

CH₄ methane profile/burden product

(a, b, c) retrieve physical parameters a, b, c

(update a, b, c) update previously retrieved parameters a,b,c